

## ERRATUM

DOI: 10.14744/AnatolJCardiol.2024.3562

The article by Erol et al., published in the May issue [Anatol J Cardiol. 2024 May 27;28(8):393–8], titled “Agreement Between Transthoracic Echocardiography and Computed Tomography Pulmonary Angiography for Detection of Right Ventricular Dysfunction in Pulmonary Embolism,” contains an error that was brought to our attention through a Letter to the Editor.

Specifically, certain values listed in Table 5 under the “Positive Predictive Value” section were found to be incorrect. The authors have since provided the correct values, which have been reviewed and approved by the Editorial Board. We sincerely apologize to our readers for this error and any confusion it may have caused.

The updated table is provided below. The article can be accessed via the following link: <https://anatoljcardiol.com/storage/upload/pdfs/1722327195-en.pdf>

**Table 5. Performance of CTPA and TTE on prediction of adverse outcomes**

	<b>Sensitivity</b>	<b>Specificity</b>	<b>Positive Predictive Value</b>	<b>Negative Predictive Value</b>
RV dysfunction on TTE	100%	43%	55%	100%
RV dysfunction on CTPA	100%	28%	41%	100%

CTPA, computed tomography pulmonary angiography;  
TTE, transthoracic echocardiography.

