

Echocardiographic imaging of saccular aneurysm in the left main coronary artery

A 59-year-old woman presented with a recent onset of dyspnea and chest pain. Past medical history and cardiovascular examination were completely normal. Electrocardiography demonstrated sinus rhythm with left bundle branch block pattern. Transthoracic echocardiography (TTE) revealed a slightly reduced ejection fraction with hypokinetic septal and anterior walls. Parasternal short-axis imaging displayed a small saccular echo-free space associated with the aorta, and no color flow turbulence was observed with color flow Doppler imaging (Fig. 1a). Subsequently, two-dimensional transesophageal echocardiography (2D-TEE) revealed a normal long-axis view and similar short-axis findings to transthoracic imaging. Real-time 3D-TEE (RT-3D-TEE) provided better imaging and indicated that the saccular body was a round-shaped small aneurysm that was relevant to the left main coronary artery (LMCA) take-off location (Fig. 1b; Video 1).

Aortography depicted a round-shaped saccular aneurysm, 8×9 mm in size, originating from the proximal LMCA that was approximately 2–3 mm next to the LMCA ostium (Fig. 2a; Video 2). Coronary angiography revealed an aneurysm associated with proximal LMCA (Fig. 2b; Video 3).

Although TTE can provide valuable information regarding the diagnosis of coronary aneurysm in pediatric patients with Kawasaki disease, its validity is limited in adults because of the declining image quality.




Figure 1. (a) 2D Transthoracic echocardiography parasternal short-axis image, from the slightly higher level of the aortic valve, displaying small saccular echo-free space (white arrows). (b) Real-time 3D transesophageal echocardiography showing small, round-shaped, proximal LMCA segment originated aneurysm (red arrows) in the diastolic short-axis image.

*represents. Ao- aorta; LA- left atrium; RA- right atrium; RV- right ventricle; LMCA- left main coronary artery; PA- pulmonary artery




Figure 2. (a) A. Aortography showing a round-shaped, proximal LMCA-originated saccular aneurysm, 8×9 mm in size (Black arrows). (b) Coronary angiography showing proximal LMCA aneurysm (black arrows) and non-significant lesions in the coronary arteries.

*LMCA. Ao- aorta; Cx- circumflex artery; LAD- left anterior descending artery

Nevertheless, proximal segments of the coronary arteries can be assessed with TEE. RT-3D-TEE can provide even better information regarding the location, size, shape, and relation to adjacent tissue in LMCA aneurysm, as in our case.

In this report, we presented an isolated saccular LMCA aneurysm diagnosed by echocardiography, which is rarely encountered in coronary angiography.

Hatem Ari, Fatih Kahraman, Fatih Aksoy, İsmail Barkın Işık
Department of Cardiology, Faculty of Medicine, Süleyman
Demirel University; Isparta-Turkey

Video 1. Real-time 3D transesophageal echocardiography showing small, saccular, proximal LMCA aneurysm at the level of the sinotubular junction

Video 2. Aortography showing saccular aneurysm originating from the proximal LMCA

Video 3. Coronary angiography showing proximal LMCA aneurysm and normal coronary arteries

Address for Correspondence: Dr. Hatem Ari

Araştırma ve Uygulama Hastanesi, Süleyman Demirel Üniversitesi Doğu

Yerleşkesi 32260, Cunur, Isparta-Türkiye

Phone: +90 246 211 20 00

Fax: +90 246 211 28 30

E-mail: hatem_ari@hotmail.com

©Copyright 2015 by Turkish Society of Cardiology - Available online at www.anatoljcardiol.com

DOI:10.5152/AnatolJCardiol.2015.6385

Three-dimensional imaging of left ventricular hemangioma

A 30-year-old asymptomatic man with chronic hepatitis B was evaluated for an incidentally visualized intracardiac mass on abdominal ultrasonography. Two-dimensional (2D) transthoracic echocardiography (TTE) revealed a structurally normal heart except for the presence of homogeneously echodense, round, mobile mass (2.0×2.8 cm) within the left ventricular cavity (Fig. 1a, Video 1a). No flow abnormalities were detected with Doppler analysis. On three-dimensional (3D) TTE, the mass appeared to be attached to the interventricular septum (Fig. 1b, Video 1b). Transesophageal echocardiography (TEE) was performed for superior visualization of anatomical details of the mass. On 2D TEE imaging with xPlane, the mass was well-demarcated and attached to the interventricular septum with a faintly visible stalk, and there was no infiltration of the myocardial wall (Fig. 2a, Video 2a). On real-time 3D TEE, the surface




Figure 1. (a) Apical five-chamber view showing a homogeneously echodense, round, mobile mass within the left ventricular cavity. (b) Real-time three-dimensional transthoracic echocardiography showing a left ventricular mass and interventricular septum.