

An interesting steal case 🎬

A 65-year-old man applied to our hospital with efor angina that have progressive raised for last 6 months. He had undergone coronary bypass operation three years ago. Also he has diabetes and hypertension. Owing to his typical symptoms a coronary angiogram was performed. In his coronary angiogram left anterior descending arter (LAD) was totally occluded at mid-level, circumflex and right coronary arteries (RCA) and aort-obtus marginalis (OM), saphenous graft were open (Fig. 1). A retrograd flow was seen between aort- diagonal grafts to left subclavian artery (LSA) through to left internal mammarian artery (LIMA) (Fig. 2A-C, Video 1, 2). Attempts was failed to demonstrate LIMA due to total occlusion of LSA (Fig. 2D, Video 3). LIMA originates from LSA. In case of LIMA graft an occlusion or severe stenosis of LSA




Figure 1. Coronary angiography of the patient (1A, 1B Left system 1C RCA, 1D Aort- OM)




Figure 2. LIMA- LAD connection and absence of distal LAD flow, (2A Retrograd flow between Aort- diagonal to LSA 2B-C. Total occlusion of LSA 2D)

causes retrograd flow from coronary arteries to left upper extremity as a consequence myocardial ischemia. Mostly retrograd flow comes from native LAD through to LIMA. In our case a saphenous graft provides connection between ascending aorta to LSA by using diagonal artery, LAD and LIMA. We consider diffuse illness of distal LAD may support that retrograd flow. We suggest that checking LSA and LIMA before the bypass operation could provide that undesirable results.

Kamil Gülşen, Levent Cerit, İrfan Şahin¹, Berk Özkaynak², Ertuğrul Okuyan¹

Department of Cardiology, Faculty of Medicine, Near East University; Mersin-Turkey

Clinic of ¹Cardiology and ²Cardiovascular Surgery, Bağcılar Education and Research Hospital; Istanbul -Turkey

Video 1. Aort- diagonal saphenous graft and retrograd flow through to LIMA.

Video 2. Filling of subclavian artery by retrograd flow.

Video 3. Totally occlusion at ostium of left subclavian artery.

Address for Correspondence: Dr. Kamil Gülşen,

Yakın Doğu Üniversitesi, Kalp Merkezi

PB 99138 Nicosia Mersin-Türkiye

Phone: +90 392 459 10 00

Fax: +90 392 223 64 61

E-mail: kamilgulsen2000@yahoo.com

Available Online Date: 06.05.2014

©Copyright 2014 by Turkish Society of Cardiology - Available online at www.anakarder.com

DOI:10.5152/akd.2014.5315

An unusual right ventricle outflow obstruction in adult 🎬

A 64-years old man who had been experiencing worsening dyspnea and atypical chest pain in the past 6 months, was admitted to our department. Upon arrival, physical examination revealed a 2-3/6 systolic murmur heard best over the left second intercostal space. Transthoracic echocardiography was promptly performed showing a thin-walled outpouching at level of right sinus of Valsalva, sized 45 x 50 mm (Fig. 1A). In parasternal short axis view, this appeared completely obliterate right ventricular outflow tract (RVOT) and color Doppler showed turbulent flow in this site; pulmonary valve was not well visualized (Fig. 1B, Video 1). Continuous wave Doppler study confirmed the obstruction with a peak velocity of 3 m/sec and peak gradient of 36 mm Hg (Fig. 1C). A contrastographic study with Sonovue® allowed to exclude a communication between the outpouching and the right ventricle (Video 2, 3). According to the anatomical site and echocardiographic findings the diagnosis of unruptured right sinus of valsalva aneurysm was made. To better define its relationship with nearest cardiac structures and to plane a correct surgical strategy, computed tomography (CT) was performed, confirming the presence of the aneurysm and the RVOT obstruction below pulmonary valve (Fig. 1D-F). The patient underwent surgical intervention; the aneurysm was removed with subsequent complete relief of the symptoms during the postoperative period.

Andrea Madeo, Paolo Giuaeppe Pino, Amedeo Pergolini,

Claudia Chialastri, Vitaliano Buffa

Department of Cardiovascular Science, "S. Camillo Forlanini" Hospital; Rome-Italy