

Reply to Letter to the Editor: "Tricuspid Regurgitation Worsening After Pericardiectomy in Tuberculosis Constrictive Pericarditis: An Overlooked Prognostic Concern"

To the Editor,

The authors¹ are thanked for their interest and valuable comments on the article "Tricuspid Regurgitation in Tuberculous Constrictive Pericarditis Underwent Pericardiectomy."² Tuberculosis infection is the main cause of constrictive pericarditis (CP) in developing countries, accounting for approximately 38%-83% of all CP cases. Currently, pericardiectomy is the only effective treatment for CP.³ Studies have shown that both CP combined with tricuspid regurgitation (TR) and worsening TR due to pericardiectomy reduce long-term survival. This phenomenon that deserves attention in the preoperative evaluation by surgeons.⁴⁻⁷

There is no consensus on the mechanism of coexisting TR in CP. It is believed that TR is secondary to the structural and functional remodeling of the patient's right atrium and right ventricle during the pathological processes of CP and that the patient's combination of underlying diseases, such as arrhythmia, pacemaker implantation, and chronic lung disease, is an important influencing factor. In tuberculous CP, the presence of inflammatory factors that damage the valvular apparatus deserves further research and discussion.

In CP, worsening TR is associated with right ventricular systolic dysfunction. Meanwhile, impaired right ventricular systolic function complicates the treatment of pericardiectomy and reduces long-term patient survival. This makes a detailed assessment of right ventricular morphology and function particularly important.

Currently, echocardiographic assessment of right ventricular systolic function is mainly based on right ventricular fractional area change, tricuspid annular plane systolic excursion, and right ventricular S' tissue Doppler velocity, as mentioned in the article, but it can be considered using speckle tracking imaging, 3-dimensional echocardiography or even magnetic resonance imaging to assess right ventricular morphology, local myocardial function, and degree of fibrosis in detail.

Thank you again for agreeing with the viewpoint of the article. For tuberculosis CP, we will prospectively study the patients through various aspects, including clinical, blood biochemical, multimodal imaging, and prognostic aspects, to understand the mechanism of worsening TR and the factors and extent of right ventricular myocardial injury. This will provide more information for surgical strategies and improve the long-term prognosis of patients.

Declaration of Interests: The authors have no conflicts of interest to declare.

Funding: This work was supported by Sichuan Natural Science Foundation (24NSFC0262).



Copyright©Author(s) - Available online at anatoljcardiol.com.
Content of this journal is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License.

LETTER TO THE EDITOR REPLY

Shuzhen Wang¹

Jing Guo¹

Kunyue Tan¹

Lijuan Zhang¹

Shuangshuang Yan¹

Li Zhang¹

Wei Wang²

Qiuyi Cai³

Feng Xiong¹

¹Department of Cardiology, The Third People's Hospital of Chengdu, The Affiliated Hospital of Southwest Jiaotong University, Chengdu, China

²Department of Cardiovascular Surgery, The Third People's Hospital of Chengdu, The Affiliated Hospital of Southwest Jiaotong University, Chengdu, China

³Department of Radiology, The Third People's Hospital of Chengdu, The Affiliated Hospital of Southwest Jiaotong University, Chengdu, China

Corresponding author:

Feng Xiong

✉ xiong.feng05@163.com

Cite this article as: Wang S, Guo J, Tan K, et al. Reply to letter to the editor: "tricuspid regurgitation worsening after pericardiectomy in tuberculosis constrictive pericarditis: an overlooked prognostic concern." *Anatol J Cardiol.* 2025;29(10):592-593.

DOI:10.14744/AnatolJCardiol.2025.5469

REFERENCES

1. Demirçelik B. Tricuspid regurgitation worsening after pericardiectomy in tuberculosis constrictive pericarditis: an overlooked prognostic concern. *Anatol J Cardiol.* 2025;29(10):590-591.
2. Wang SZ, Guo J, Tan KY, et al. Tricuspid regurgitation in tuberculous constrictive pericarditis underwent pericardiectomy. *Anatol J Cardiol.* 2025;29(5):222-227. [\[CrossRef\]](#)
3. Mayosi BM, Burgess LJ, Doubell AF. Tuberculous pericarditis. *Circulation.* 2005;112(23):3608-3616. [\[CrossRef\]](#)
4. Calderon-Rojas R, Greason KL, King KS, et al. Tricuspid valve regurgitation in patients undergoing pericardiectomy for constrictive pericarditis. *Semin Thorac Cardiovasc Surg.* 2020;32(4):721-728. [\[CrossRef\]](#)
5. Tabucanon RS, Wang TKM, Chetrit M, et al. Worsened tricuspid regurgitation following pericardiectomy for constrictive pericarditis. *Circ Cardiovasc Imaging.* 2021;14(10):e012948. [\[CrossRef\]](#)
6. Homsy M, Mahenthiran J, Vaz D, Sawada SG. Reduced right ventricular systolic function in constrictive pericarditis indicates myocardial involvement and persistent right ventricular dysfunction and symptoms after pericardiectomy. *J Am Soc Echocardiogr.* 2007;20(12):1417.e1-1417.e714177. [\[CrossRef\]](#)
7. Choudhry MW, Homsy M, Mastouri R, Feigenbaum H, Sawada SG. Prevalence and prognostic value of right ventricular systolic dysfunction in patients with constrictive pericarditis who underwent pericardiectomy. *Am J Cardiol.* 2015;116(3):469-473. [\[CrossRef\]](#)