

<sup>1</sup>Department of Cardiology, Faculty of Medicine, Selçuk University; Konya-Turkey

<sup>2</sup>Cardiology Clinics, Afyonkarahisar Dinar State Hospital; Afyonkarahisar-Turkey

## References

1. Gürses KM, Yalçın MU, Koçyiğit D, Beşler MS, Canpınar H, Evranos B, et al. The association between serum angiogenin and osteopontin levels and coronary collateral circulation in patients with chronic total occlusion. *Anatol J Cardiol* 2019; 22: 77-84.
2. Tello-Montoliu A, Marín F, Patel J, Roldán V, Mainar L, Vicente V, et al. Plasma angiogenin levels in acute coronary syndromes: implications for prognosis. *Eur Heart J* 2007; 28: 3006-11.
3. Huang SD, Lu FL, Xu XY, Liu XH, Zhao XX, Zhao BZ, et al. Transplantation of angiogenin-overexpressing mesenchymal stem cells synergistically augments cardiac function in a porcine model of chronic ischemia. *J Thorac Cardiovasc Surg* 2006; 132: 1329-38.
4. Kishimoto K, Liu S, Tsuji T, Olson KA, Hu GF. Endogenous angiogenin in endothelial cells is a general requirement for cell proliferation and angiogenesis. *Oncogene* 2005; 24: 445-56.

**Address for Correspondence:** Dr. Muhammed Ulvi Yalçın, Selçuk Üniversitesi Tıp Fakültesi, Kardiyoloji Anabilim Dalı, Konya-Türkiye  
Phone: +90 332 221 00 00  
Fax: +90 332 323 67 23  
E-mail: ulviyalcin@gmail.com  
©Copyright 2019 by Turkish Society of Cardiology - Available online at [www.anatoljcardiol.com](http://www.anatoljcardiol.com)

## Methodological problems in the measurement of interleukin-33 concentrations in patients with heart failure with a reduced ejection fraction

To the Editor,

We have read manuscript entitled "The concentration of interleukin-33 (IL-33) in heart failure with reduced ejection fraction. *Anatol J Cardiol* 2019; 21: 305-13" with great interest (1). The study included 155 patients with heart failure and 60 healthy individuals to compare IL-33 levels (1). The authors pointed out that blood samples collected from patients on admission were stored at -80 °C until further use; however, they did not mention about blood sample storage time comparison between the patient and the control group. Storage time for blood samples may affect IL-33 concentration measurement and change results; for example, variable cytokine concentration levels were seen with different blood storage durations in the previous study (2).

The authors used the legend max human IL-33 kit manufacturer system as seen in the standard range of 15.6–1000 pg/mL, and the limit of detection is 4.14 pg/mL; it is not clear how values below this level are measured (3). However, the authors measured a healthy human concentration of 16.91 (0–81) pg/mL in Figure 1a, which we consider as an important problem as to how the kit measured levels below the limit of detection. Values between the limit of quantification and the limit of detection should be given qualitatively and calculated separately. Furthermore, we think that statistical measurement and interpretations made without value and corrections may affect the results.

Figure 1b is given as an interquartile range in figure legends, but that is not found in the text or figure. No numerical value was given in the text or figure for Figure 1c. In addition, the median was given in Figure 1d, and it was observed that the median value was below the limit of quantification.

Therefore, we think that the interpretations made on IL-33 in Figure 1a–1d are insufficient due to the methodological approach, and that the statistical comments about the values will seriously reduce the reliability. Furthermore, how values <4.14 are measured raises considerable doubt as they are below the minimum measurable limit.

The aim of the present study was to investigate the effect of cytokine expression on the course of the disease in patients with heart failure. However, case-control studies do not provide information about the course of the disease. These studies can only comment on the instant situation. To do this, we think that evaluation with longitudinal cohort studies would be a more appropriate method.

 Berhan Keskin,  İsmail Balaban,  Seda Tanyeri,  Özgür Yaşar Akbal,  Ali Karagöz

Department of Cardiology, Koşuyolu Kartal Heart Training and Research Hospital; Istanbul-Turkey

## References

1. Segiet OA, Romuk E, Nowalany-Kozielska E, Wojciechowska C, Piecuch A, Wojnicz R. The concentration of interleukin-33 in heart failure with reduced ejection fraction. *Anatol J Cardiol* 2019; 21: 305-13.
2. Jackman RP, Utter GH, Heitman JW, Hirschhorn DF, Law JP, Geffer N, et al. Effects of blood sample age at time of separation on measured cytokine concentrations in human plasma. *Clin Vaccine Immunol* 2011; 18: 318-26. [CrossRef]
3. <https://www.biolegend.com/en-us/products/legend-max-human-il-33-elisa-kit-with-pre-coated-plates-6433>.

**Address for Correspondence:** Dr. Berhan Keskin, Kartal Koşuyolu Yüksek İhtisas Eğitim ve Araştırma Hastanesi, Kardiyoloji Bölümü, Denizer Cad. Cevizli Kavşağı No: 2, Kartal, İstanbul-Türkiye  
Phone: +90 537 977 67 36  
E-mail: bekeskin@ku.edu.tr

©Copyright 2019 by Turkish Society of Cardiology - Available online at [www.anatoljcardiol.com](http://www.anatoljcardiol.com)  
DOI:10.14744/AnatolJCardiol.2019.59387

