

Is a Single Nitroglycerin Study Sufficient to Say That It Is Safe Not to Give Nitroglycerin Before Radial Intervention?

To the Editor,

We read with great interest the article titled "Impact of Nitroglycerine Injection on Radial Artery Outcomes in Distal Transradial Access" written by Karaca et al¹ and published in *Anatol J Cardiol* 2025; 29(3): 139-144. This study comes across as a well-designed study on the benefit/disuse of nitroglycerin injection before radial artery intervention. However, there are some questions that we would like to ask the authors politely. It is stated in the study that the study was conducted retrospectively. It is seen that the patients who applied were accepted as elective or emergency cases. However, are the troponin values given in Table 1 within the normal range according to the applications? Or we request that the normal-high value of troponin be stated in the article. We would also like to ask why the patients included in the study did not receive nitroglycerin before radial intervention, although there are studies recommending nitroglycerin.² Also, 1 study³ found that patients with radial artery diameters and hyperlipidemia may cause radial artery occlusion after or during the procedure. May we ask if the authors' study contains data on this?

Declaration of Interests: The authors have no conflicts of interest to declare.

Funding: The authors declare that this study received no financial support.

REFERENCES

1. Karaca M, Dağdeviren B. Impact of nitroglycerine injection on radial artery outcomes in distal transradial access. *Anatol J Cardiol*. 2025;29(3):139-144. [CrossRef]
2. Kim SH, Kim EJ, Cheon WS, et al. Comparative study of nicorandil and a spasmolytic cocktail in preventing radial artery spasm during transradial coronary angiography. *Int J Cardiol*. 2007;120(3):325-330. [CrossRef]
3. Avdikos G, Karatasakis A, Tsoumeleas A, Lazaris E, Ziakas A, Koutouzis M. Radial artery occlusion after transradial coronary catheterization. *Cardiovasc Diagn Ther*. 2017;7(3):305-316. [CrossRef]

LETTER TO THE EDITOR

Sefa Erdi Ömür 

Çağrı Zorlu 

Department of Cardiology, Faculty of
Medicine, Tokat Gaziosmanpaşa
University, Tokat, Türkiye

Corresponding author:

Sefa Erdi Ömür
✉ sefaerdi61@gmail.com

Cite this article as: Ömür SE, Zorlu Ç. Is a single nitroglycerin study sufficient to say that it is safe not to give nitroglycerin before radial intervention? *Anatol J Cardiol*. 2025;XX(X):1.



Copyright©Author(s) - Available online at anatoljcardiol.com.
Content of this journal is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License.

DOI:10.14744/AnatolJCardiol.2025.5528