

Mitral regurgitation secondary to mitral anterior leaflet rupture after mitral valvuloplasty



Mitral valvüloplasti sonrası mitral ön kapak rüptürüne sekonder olarak gelişen mitral yetmezlik

A 32-year-old female patient was admitted to our outpatient clinic for routine checkup. In her history, she underwent percutaneous mitral balloon valvuloplasty due to mitral stenosis three months ago. On physical examination, an apical 2/6 systolic murmur was heard. Electrocardiography showed normal sinus rhythm. Two-dimensional transthoracic echocardiography revealed eccentric mitral regurgitation, which was not present previously. To clarify this pathology, we performed two- and three-dimensional transesophageal echocardiography (2-D and 3-D TEE). 2-D TEE mid-esophageal view demonstrated a cleft on the mitral anterior leaflet (Fig. 1A arrow and Video 1A. See corresponding video/movie images at www.anakarder.com). Color Doppler echocardiography showed a mitral regurgitation resulting from this cleft (Fig.1B and Video 1B. See corresponding video/movie images at www.anakarder.com). 3-D color full volume TEE confirmed mitral regurgitation at the anterior mitral leaflet (Fig. 1C and Video 1C. See corresponding video/movie images at www.anakarder.com). 3D Live Zoom acquisition revealed the cleft at A2 scallop (Fig. 1D arrow and Video 1D. See corresponding video/movie images at www.anakarder.com).

In patients with mitral regurgitation, three-dimensional echocardiography may provide the etiology of the mitral regurgitation and the correct localization of the pathology and guarantee optimal surgical guidance.




Figure 1. 2-D TEE views showing a perforation on the anterior mitral valve (A), color Doppler 2-D TEE views revealing a mitral regurgitation resulting from this perforation (B), 3-D color full volume TEE confirmed mitral regurgitation at the anterior mitral leaflet (C) and 3D Live Zoom acquisition revealed the cleft at A2 scallop

2-D - two-dimensional, 3-D - three-dimensional, TEE - transesophageal echocardiography

Video 1A. Two-dimensional transesophageal echocardiography (2-D TEE) showing a perforation on the anterior mitral valve

Video 1B. Color Doppler 2-D TEE revealing a mitral regurgitation resulting from this perforation

2-D TEE - two-dimensional transesophageal echocardiography

Video 1C. 3-D color full volume TEE confirmed mitral regurgitation at the anterior mitral leaflet

3-D - three-dimensional, TEE - transesophageal echocardiography

Video 1D. 3D Live Zoom acquisition revealed the cleft at A2 scallop

3-D - three-dimensional

Sait Demirkol, Murat Ünlü¹, Şevket Balta, Atıla İyisoy
Department of Cardiology, Faculty of Medicine, Gülhane Military Medical Academy, Ankara-Turkey
¹Department of Cardiology, Beytepe Military Hospital, Ankara-Turkey

Address for Correspondence/Yazışma Adresi: Dr. Sait Demirkol
 Gülhane Askeri Tıp Akademisi, Kardiyoloji Bölümü, Tevfik Sağlık Cad. 06018
 Etlik, Ankara-Türkiye
 Phone: +90 312 304 42 81 Fax: +90 312 304 42 50
 E-mail: saitdemirkol@yahoo.com

Available Online Date/Çevrimiçi Yayın Tarihi: 18.09.2012

©Telif Hakkı 2012 AVES Yayıncılık Ltd. Şti. - Makale metnine www.anakarder.com web sayfasından ulaşılabilir.

©Copyright 2012 by AVES Yayıncılık Ltd. - Available on-line at www.anakarder.com
 doi:10.5152/akd.2012.234

Incidental diagnosis of an aneurysm of the mitral valve posterior leaflet



Mitral kapak arka yaprak anevrizmasının rastlantısal tanısı

Mitral valve aneurysms are also most frequently associated with endocarditis, but in rare cases of mitral aneurysms are associated with connective tissue diseases or congenital malformation. Transesophageal echocardiography is a more sensitive tool than transthoracic echocardiography to identify aneurysm.

A 35-year-old male patient was admitted to our clinic with progressive shortness of breath lasting for one year. He had no history of heart disease, endocarditis and connective tissue disease. On his physical examination, blood pressure and heart rate were 115/60 mmHg and 80 bpm respectively. Heart and respiratory auscultation findings were normal except 4/6 systolic murmur, which was heard maximally at the apex. Electrocardiogram showed sinus rhythm. Laboratory findings did not suggest of infection, blood cultures were negative and he had not fever history. Transthoracic echocardiography showed severe mitral regurgitation and a localized bulge of the mitral posterior leaflet toward the left atrium with systolic expansion and diastolic collapse. Transesophageal echocardiography revealed a thin-walled, saccular 0.7x1.0 cm mitral valve aneurysm on the left atrial surface, expanding and decompressing during cardiac cycle (Fig. 1A, Video 1. See corresponding video/movie images at www.anakarder.com). Color Doppler imaging confirmed severe mitral regurgitation and communication of aneurysm with left ventricular cavity (Fig. 1B, Video 2. See corresponding video/movie images at www.anakarder.com). Mitral valve aneurysms were confirmed at operation and valve repair was performed. After operation, echocardiography showed no significantly mitral regurgitation. The patient was discharged uneventfully.




Figure 1. Transesophageal 2-chamber echocardiographic view of an aneurysm of the mitral posterior leaflet (A) and color-flow Doppler image demonstrating a severe mitral regurgitation (B)

LA- left atrium, LV-left ventricle, RA- right atrium, RV- right ventricle.

Video 1. Transesophageal echocardiography views of a saccular mitral valve aneurysm on the left atrial surface, expanding and decompressing during cardiac cycle

Video 2. Color Doppler imaging of a severe mitral regurgitation and communication of aneurysm with left ventricular cavity

Turgay Işık, Mustafa Kurt¹, Erkan Ayhan, İbrahim Halil Tanboğa¹
Department of Cardiology, Faculty of Medicine, Balıkesir University, Balıkesir-Turkey
¹Clinic of Cardiology, Erzurum Education and Research Hospital, Erzurum-Turkey

Address for Correspondence/Yazışma Adresi: Dr.Turgay Işık
Balıkesir Üniversitesi Tıp Fakültesi, Kardiyoloji Anabilim Dalı, Çağış Kampüsü,
Balıkesir-Türkiye
Phone: +90 266 612 14 55 Fax: +90 266 612 14 59
E-mail: isikturgay@yahoo.com

Available Online Date/Çevrimiçi Yayın Tarihi: 18.09.2012

©Telif Hakkı 2012 AVES Yayıncılık Ltd. Şti. - Makale metnine www.anakarder.com web sayfasından ulaşılabilir.

©Copyright 2012 by AVES Yayıncılık Ltd. - Available on-line at www.anakarder.com
doi:10.5152/akd.2012.235

Incidentally diagnosed pseudoaneurysm of mitral-aortic intervalvular fibrosa



Rastlantısal tanı konulmuş mitral-aortik intervalvular fibroza psödoanevrizması

Pseudoaneurysm of mitral-aortic intervalvular fibrosa (P-MAIVF) is defined as a pseudoaneurysm at interannular zone between mitral and aortic valves. Communication of the cavity with the ventricular outflow tract and the pulsatility of the cavity during cardiac cycle are features differentiating pseudoaneurysms from ring abscesses. Most common associated reasons include endocarditis and aortic valve surgery.

A 77-year-old man with mechanical aortic valve replacement due to aortic stenosis underwent transthoracic echocardiography (TTE) before non-cardiac surgery. TTE revealed an echolucent area adjacent to aortic valve, raising suspicion of a P-MAIVF (Fig.1, Video 1. See corresponding video/movie images at www.anakarder.com). He had no chest pain or dyspnea and any sign of infective endocarditis or prosthetic valve dysfunction. Transesophageal echocardiography (TEE) showed an echolucent area measuring 15 mm in its widest dimension, which was communicating with left ventricular outflow tract (LVOT). This cavity was



Figure 1. Transthoracic echocardiography views showing echolucent area adjacent to the aortic valve (arrow)

AO - ascending aorta, LA - left atrium, LV - left ventricle




Figure 2. Transesophageal echocardiography views showing a pseudoaneurysm in the mitral-aortic intervalvular fibrosa (arrow)

AO - ascending aorta, LA - left atrium, LV - left ventricle